

Unitarity and universality with kaon physics at KLOE

Erika De Lucia for the KLOE Collaboration *

Laboratori Nazionali di Frascati dell'INFN, Roma, Italia.

All relevant inputs for the extraction of the CKM matrix element V_{us} from K_L , K_S and $K^\pm$ decays have been measured at KLOE. From a global fit using only KLOE results, but K_S lifetime, a value of $|V_{us}|f_+(0) = 0.2157 \pm 0.0006$ is obtained, where $f_+(0)$ is the form factor parametrizing the hadronic matrix element evaluated at zero momentum transfer. Comparison of the values of $f_+(0)V_{us}$ for K_{e3} and $K_{\mu 3}$ modes provides a test of lepton universality at 0.8% level of accuracy. The value $V_{us}/V_{ud} = 0.2323(15)$ has been obtained from the ratio $\Gamma(K \rightarrow \mu\nu)/\Gamma(\pi \rightarrow \mu\nu)$ using KLOE measurement of $\text{BR}(K^\pm \rightarrow \mu^\pm\nu)$ and lattice calculation of the ratio of decay constants f_K/f_π . These results, together with $V_{ud} = 0.97418(26)$, are compatible at 0.6σ level with CKM matrix unitarity. The universality of lepton and quark weak couplings can be tested and constraints on new physics extensions of the Standard Model can be set using these very precise measurements from kaon decays.

1. INTRODUCTION

Precise measurements of semileptonic kaon decay rates provide the measurement of the V_{us} element of the CKM mixing matrix and information about lepton universality. Helicity-suppressed leptonic kaon decays provide an independent measurement of $|V_{us}|^2/|V_{ud}|^2$, through the ratio $\Gamma(K \rightarrow \mu\nu)/\Gamma(\pi \rightarrow \mu\nu)$. These measurements, together with the result of $|V_{ud}|$ from nuclear β transitions, provide the most precise test of the unitarity of the CKM mixing matrix through the relation $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$. Unitarity can also be interpreted as a test of the universality of lepton and quark weak couplings, testing the relation $G_F^2 = G_{CKM}^2 = G_F^2(V_{ud}^2 + V_{us}^2 + V_{ub}^2)$, with G_F the Fermi coupling constant from the muon decay. Bounds on new physics extensions of the standard model can be set using the ratio of the V_{us} values obtained from helicity-suppressed $K_{\ell 2}$ decays and helicity-allowed $K_{\ell 3}$ decays. The kaon semileptonic decay rate is given by:

$$\Gamma(K_{l3}) = \frac{C_K^2 G_F^2 M_K^5}{192\pi^3} S_{EW} |V_{us}|^2 |f_+(0)|^2 I_{K,l}(\lambda) (1 + 2\Delta_K^{SU(2)} + 2\Delta_{K,l}^{EM}) \quad (1)$$

where $K = K^0, K^\pm$, $l = e, \mu$ and C_K is a Clebsch-Gordan coefficient, equal to $1/2$ and 1 for $K^\pm$ and K^0 , respectively. The decay width $\Gamma(K_{l3})$ is experimentally determined by measuring the kaon lifetime and the semileptonic BRs totally inclusive of radiation. The theoretical inputs are: the universal short-distance electroweak correction $S_{EW} = 1.0232$, the $SU(2)$ -breaking $\Delta_K^{SU(2)}$ and the long-distance electromagnetic corrections $\Delta_{K,l}^{EM}$, which depend on the kaon charge and on the lepton flavor, and the form factor $f_+(0) \equiv f_+^{K^0\pi^-}(0)$ parametrizing the hadronic matrix element of the $K \rightarrow \pi$ transition, evaluated at zero momentum transfer and for neutral kaons. The form factor dependence on the momentum transfer can be described by one or more slope parameters λ , measured from the decay spectra, and enters in the phase space integral $I_{K,l}(\lambda)$.

The KLOE experiment at the Frascati ϕ -factory DAΦNE can measure all the relevant inputs to extract V_{us} from K_{l3} decay rates of both charged and neutral kaons: BRs, lifetimes and form factors. In section 2 an overview of the measurements for K_L , K_S and $K^\pm$ will be presented, followed by the extraction of V_{us} in section 3 and bounds on new physics beyond the standard model in section 4.

*F. Ambrosino, A. Antonelli, M. Antonelli, F. Archilli, C. Bacci, P. Beltrame, G. Bencivenni, S. Bertolucci, C. Bini, C. Bloise, S. Bocchetta, F. Bossi, P. Branchini, P. Campana, G. Capon, T. Capussela, F. Ceradini, S. Chi, G. Chieffari, P. Ciambriano, E. De Lucia, A. De Santis, P. De Simone, G. De Zorzi, A. Denig, A. Di Domenico, C. Di Donato, B. Di Micco, A. Doria, M. Dreucci, G. Felici, A. Ferrari, M. L. Ferrer, S. Fiore, C. Forti, P. Franzini, C. Gatti, P. Gauzzi, S. Giovannella, E. Gorini, E. Graziani, W. Kluge, V. Kulikov, F. Lacava, G. Lanfranchi, J. Lee-Franzini, D. Leone, M. Martini, P. Massarotti, W. Mei, S. Meola, S. Miscetti, M. Moulson, S. Müller, F. Murtas, M. Napolitano, F. Nguyen, M. Palutan, E. Pasqualucci, A. Passeri, V. Patera, F. Peretto, M. Primavera, P. Santangelo, G. Saracino, B. Sciascia, A. Sciubba, A. Sibidanov, T. Spadaro, M. Testa, L. Tortora, P. Valente, G. Venanzoni, R. Versaci, G. Xu.

2. BRANCHING RATIOS, LIFETIMES AND FORM FACTOR SLOPES

2.1. K_L decays

The absolute BRs for the four main K_L decay channels have been measured from a sample of 1.3×10^7 $\phi \rightarrow K_S K_L$ events tagged by $K_S \rightarrow \pi^+ \pi^-$ decays. The results depend on the K_L lifetime τ_L through the geometrical acceptance of the apparatus. Using as reference value $\tau_L = 51.54$ ns we get $\text{BR}(K_{Le3}) = 0.4049(21)$, $\text{BR}(K_{L\mu3}) = 0.2726(16)$, $\text{BR}(3\pi^0) = 0.2018(24)$, and $\text{BR}(\pi^+ \pi^- \pi^0) = 0.1276(15)$ [1]. An independent measurement, $\tau_L = 50.92(30)$ ns [2], is obtained from the fit of the proper decay time distribution for $K_L \rightarrow 3\pi^0$ events. K_L tagged decays have been used for the measurement of the form factors for both K_{e3} and $K_{\mu3}$ decays. For K_{e3} decays the fit of the t/m_π^2 distribution, with t the squared momentum transfer, is sensitive to both linear and quadratic terms of the power expansion of the form factor and agreement has been found between the results obtained with quadratic and pole parametrization. With the quadratic parametrization we get: $\lambda'_+ = 25.5(1.8) \times 10^{-3}$ and $\lambda''_+ = 1.4(0.8) \times 10^{-3}$ [3]. Using $K_{\mu3}$ decays, the λ_0 slope of the scalar form factor is obtained fitting the distribution of the neutrino energy E_ν , due to the difficult π/μ identification at low energies, while λ'_+ and λ''_+ are measured from the combined fit with K_{e3} decays. The results are: $\lambda'_+ = 25.6(1.7) \times 10^{-3}$, $\lambda''_+ = 1.5(0.8) \times 10^{-3}$ and $\lambda_0 = 15.5(2.2) \times 10^{-3}$ [4]. The fit to data distributions is only sensitive to the linear term of the power expansion of the scalar form factor, due to the -99.6% correlation between the coefficients of the linear λ'_0 and quadratic terms λ''_0 . However the linear parametrization over-estimates the value of λ'_0 by about 20%. The solution is to use the dispersive representation in which vector and scalar form factors are described by two parameters only [5]: λ_+ and λ_0 . The results are $\lambda_+ = 25.7(0.6) \times 10^{-3}$ and $\lambda_0 = 14.0(2.1) \times 10^{-3}$ [4] and have been used in the evaluation of V_{us} . From a preliminary update based on a larger $K_{\mu3}$ data sample, averaged with the published values, we get: $\lambda_+ = 26.0(0.5) \times 10^{-3}$ and $\lambda_0 = 15.1(1.4) \times 10^{-3}$.

2.2. K_S decays

A ϕ -factory provides the unique opportunity of having a pure K_S -beam. Therefore, using a sample of 1.2×10^8 $\phi \rightarrow K_S K_L$ events in which the K_L is identified by its interaction in the calorimeter, we have measured $\Gamma(K_{Se3})/\Gamma(\pi^+ \pi^-) = 10.19(13) \times 10^{-4}$ [6] and $\Gamma(\pi^+ \pi^-)/\Gamma(2\pi^0) = 2.2549(54)$ [7]. These two ratios completely determine the value of K_S main BRs and allow us to measure $\text{BR}(K_{Se3}) = 7.046(91) \times 10^{-4}$ to be used in the V_{us} extraction. The value of K_S lifetime used in the present determination of $|V_{us}|f_+(0)$ is $\tau_S = 0.08958(5)$ ns, from PDG fit to CP parameters [8].

2.3. $K^\pm$ decays

The values of $\text{BR}(K_{e3}^\pm)$ and $\text{BR}(K_{\mu3}^\pm)$ have been determined separately for K^+ and K^- decays tagged by K^+ and K^- so to have four independent measurements for each BR, minimizing the systematic contribution from the tagging procedure. The background rejection uses kinematics and the signal count is extracted from a constrained likelihood fit to the distribution of the squared lepton mass evaluated by time-of-flight measurements. Using $\tau_\pm = 12.385(25)$ ns [8] to account for acceptance dependence on kaon lifetime ($\tau_\pm$), we obtain $\text{BR}(K_{e3}^\pm) = 0.04965(53)$, and $\text{BR}(K_{\mu3}^\pm) = 0.03233(39)$ [9]. The world average accuracy on $K^\pm$ lifetime $\tau_\pm$ is 0.2% [8] however the measurements' spread signals a poor consistency between the results. The KLOE experiment has measured $\tau_\pm$ with $K^\pm \rightarrow \mu^\pm \nu$ tagged kaons and fitting the distribution of the kaon proper decay time t^* evaluated with two independent methods: the kaon decay length and the kaon decay time from the time of flight of the photons from the π^0 in the final state. The first method gives $\tau_\pm = 12.364(31)(31)$ ns, the second $\tau_\pm = 12.337(30)(20)$ ns and the average is $\tau_\pm = 12.347(30)$ ns [10], in perfect agreement with PDG value. The absolute branching ratio of the $K^+ \rightarrow \pi^+ \pi^0$ (γ) decay has been measured using ~ 20 million tagged K^+ mesons. Signal counts are obtained from the fit of the distribution of the momentum of the charged decay particle in the kaon rest frame. The result, inclusive of final-state radiation, is $\text{BR}(K^+ \rightarrow \pi^+ \pi^0 (\gamma)) = 0.2065(9)$ [11], 1.3% ($\sim 2\sigma$) lower than the PDG fit [8]. We then fit the six largest $K^\pm$ BRs and the lifetime τ using our measurements of $\tau_\pm$ [10], $\text{BR}K_{\pi2}^+$ [11] $\text{BR}K_{\mu2}^+$ [12], $\text{BR}K_{13}^\pm$ [9]

$\text{BR}(K_{\mu 2}^+)$	$\text{BR}(K_{\pi 2}^+)$	$\text{BR}(\pi^\pm \pi^+ \pi^-)$	$\text{BR}(K_{e 3}^\pm)$	$\text{BR}(K_{\mu 3}^\pm)$	$\text{BR}(\pi^\pm \pi^0 \pi^0)$	$\tau_\pm (\text{ns})$
0.6376(12)	0.2071(9)	0.0553(9)	0.0498(5)	0.0324(4)	0.01765(25)	12.344(29)

Table I: Results of the fit to $K^\pm$ BRs and lifetime.

and $\text{BR} K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$ [13], with their dependence on $\tau_\pm$, together with $\text{BR}(K^\pm \rightarrow \pi^\pm \pi^+ \pi^-)$ from the PDG04 average [14], with the sum of the BRs constrained to unity. The fit result, listed in Table I, with $\chi^2/\text{ndf} = 0.59/1$ (CL=44%) shows very good consistency [11].

3. EXTRACTION OF $|V_{us}|f_+(0)$ AND $|V_{us}|$

To extract $|V_{us}|f_+(0)$ we need the $SU(2)$ -breaking [15] and the long distance EM corrections to the full inclusive decay rate [15, 16], evaluated for the first time for $K_{\mu 3}$ channels of both neutral and charged kaons [17]. The values of $|V_{us}|f_+(0)$ measured for $K_L e 3$, $K_L \mu 3$, $K_S e 3$, $K^\pm e 3$, and $K^\pm \mu 3$ decay modes are shown in Table II. The five decay modes agree well within the quoted errors and average to $|V_{us}|f_+(0) = 0.2157 \pm 0.0006$, with $\chi^2/\text{ndf} = 7.0/4$ (Prob=13%), to be compared with the world average $|V_{us}|f_+(0) = 0.2166 \pm 0.0005$ [18].

Mode	$K_L e 3$	$K_L \mu 3$	$K_S e 3$	$K^\pm e 3$	$K^\pm \mu 3$
$ V_{us} f_+(0)$	0.2155(7)	0.2167(9)	0.2153(14)	0.2152(13)	0.2132(15)

Table II: Values of $|V_{us}|f_+(0)$ extracted from K_{l3} decay rates.

Defining the ratio $r_{\mu e} = |f_+(0) V_{us}|_{\mu 3}^2 / |f_+(0) V_{us}|_{e 3}^2$ and using eq.1, we have $r_{\mu e} = g_\mu^2 / g_e^2$, with g_ℓ the coupling strength at the $W \rightarrow \ell \nu$ vertex. Lepton universality can be then tested comparing the measured value of $r_{\mu e}$ and its Standard Model prediction $r_{\mu e}^{SM} = 1$. Averaging between charged and neutral modes, we obtain $r_{\mu e} = 1.000(8)$, to be compared with what obtained from leptonic pion decays, $(r_{\mu e})_\pi = 1.0042(33)$ [19], and from leptonic τ decays $(r_{\mu e})_\tau = 1.0005(41)$ [20]. Using the determination of $|V_{us}|f_+(0)$ from K_{l3} decays and the preliminary result $f_+(0) = 0.964(5)$ from the UKQCD/RBC Collaboration [21], we get $|V_{us}| = 0.2237(13)$. Furthermore a measurement of V_{us}/V_{ud} can be obtained from a comparison of the radiative inclusive decay rates of $K^\pm \rightarrow \mu^\pm \nu(\gamma)$ and $\pi^\pm \rightarrow \mu^\pm \nu(\gamma)$ combined with a lattice calculation of f_K/f_π [22]. Using $\text{BR}(K^\pm \rightarrow \mu^\pm \nu) = 0.6366(17)$ from KLOE [12] and the preliminary lattice result $f_K/f_\pi = 1.189(7)$ from the HP/UKQCD '07 [23], we get $V_{us}/V_{ud} = 0.2323(15)$. This value can be used in a fit together with the measurements of V_{us} from K_{l3} decays and $V_{ud} = 0.97418(26)$ [24], as shown in figure 1 left.




Figure 1: Left: Results of fits to $|V_{ud}|$, $|V_{us}|$, and $|V_{us}|/|V_{ud}|$. Right: Excluded region in the $m_{H^\pm} - \tan\beta$ plane by the measurement $R_{\ell 23}$; the region excluded by $B \rightarrow \tau\nu$ is also indicated.

The result of this fit is $V_{ud} = 0.97417(26)$ and $V_{us} = 0.2249(10)$, with $\chi^2/\text{ndf} = 2.34/1$ (Prob= 13%), from which we get $1 - V_{ud}^2 + V_{us}^2 + V_{ub}^2 = 4(7) \times 10^{-4}$ compatible with unitarity at 0.6σ level. Using these results, we then

evaluate $G_{CKM} = G_F(V_{ud}^2 + V_{us}^2 + V_{ub}^2)^{1/2} = (1.16614 \pm 0.00040) \times 10^{-5} \text{ GeV}^{-2}$ which is in perfect agreement with the measurement from the muon lifetime $G_F = (1.166371 \pm 0.000006) \times 10^{-5} \text{ GeV}^{-2}$ and is competitive with the present accuracy of the measurements from tau-lepton decays and electroweak precision tests.

4. BOUNDS ON NEW PHYSICS

A particularly interesting observable is the ratio of the V_{us} values obtained from helicity-suppressed and helicity-allowed kaon modes: $R_{\ell 23} = |V_{us}(K_{\ell 2})/V_{us}(K_{\ell 3}) \times V_{ud}(0^+ \rightarrow 0^+)/V_{ud}(\pi_{\mu 2})|$, which is equal to 1 in the SM. The presence of a scalar current due to a charged Higgs H^+ exchange is expected to lower the value of $R_{\ell 23}$ [25]:

$$R_{\ell 23} = \left| 1 - \frac{m_{K^+}^2}{m_{H^+}^2} \left(1 - \frac{m_{\pi^+}^2}{m_{K^+}^2} \right) \frac{\tan^2 \beta}{1 + 0.01 \tan \beta} \right|, \quad (2)$$

with $\tan \beta$ the ratio of the two Higgs vacuum expectation values in the MSSM. To evaluate $R_{\ell 23}$, we fit our experimental data on $K_{\mu 2}$ and $K_{\ell 3}$ decays, using as external inputs the most recent lattice determinations of $f_+(0)$ [21] and f_K/f_π [23], the value of V_{ud} from [24], and $V_{ud}^2 + V_{us}(K_{l3})^2 = 1$ as a constraint. We obtain $R_{\ell 23} = 1.008 \pm 0.008$ [18]. Figure 1 right shows the region excluded at 95% CL in the charged Higgs mass m_{H^+} and $\tan \beta$ plane together with the bounds from $\text{BR}(B \rightarrow \tau \nu)$ [26]. The figure shows that Kaon physics and B physics provide complementary information.

References

- [1] F. Ambrosino et al., *Phys. Lett. B* **632**, 43 (2006)
- [2] F. Ambrosino et al., *Phys. Lett. B* **626**, 15 (2005)
- [3] F. Ambrosino et al., *Phys. Lett. B* **636**, 166 (2006)
- [4] KLOE Collaboration, F. Ambrosino et al., *JHEP* **12** (2007), 105
- [5] V. Bernard et al., *Phys. Lett. B* **638**, 166 (2006)
- [6] F. Ambrosino et al., *Phys. Lett. B* **636**, 173 (2006)
- [7] F. Ambrosino et al., *Eur. Phys. J. C* **48**, 767 (2006)
- [8] W.-M. Yao et al., *Review of Particle Physics*, *J. Phys. G* **33**, 1 (2006)
- [9] KLOE Collaboration, F. Ambrosino et al., *JHEP* **02** (2008), 098 and arXiv:0712.3841.
- [10] KLOE Collaboration, F. Ambrosino et al., *JHEP* **01** (2008), 073 and arXiv:0712.1112.
- [11] KLOE Collaboration, F. Ambrosino et al., *Phys. Lett. B* **666**, 305 (2008).
- [12] KLOE Collaboration, F. Ambrosino et al., *Phys. Lett. B* **632**, 76 (2006).
- [13] KLOE Collaboration, A. Aloisio et al., *Phys. Lett. B* **597**, 139 (2004).
- [14] S. Eidelman et al., *Phys. Lett. B* **592**, 1 (2004).
- [15] V. Cirigliano et al., *Eur. Phys. J. C* **23**, 121 (2002)
- [16] S. Descotes-Genon and B. Moussallam, *Eur. Phys. J. C* **42**, 403 (2005)
- [17] V. Cirigliano, *Proc. KAON International conference*, May 21-25, 2007 Frascati (Italy), PoS(KAON)007
- [18] KLOE Collaboration, F. Ambrosino et al., *JHEP* **04** (2008), 059
- [19] M.J. Ramsey-Musolf, S. Su and S. Tulin, *Phys. Rev. D* **76**, 095017 (2007).
- [20] M. Davier, A. Hocker and Z. Zang, *Rev. Mod. Phys.* **78**, 1043 (2006).
- [21] UKQCD/RBC Collaboration, *Preprint* arXiv:0710.5136.
- [22] W.J. Marciano, *Phys. Rev. Lett.* **93**, 231803 (2004)
- [23] HPQCD and UKQCD Collaborations, *Phys. Rev. Lett.* **100**, 062002 (2008)
- [24] I. S. Towner and J. C. Hardy *Phys. Rev. C* **77**, 025501 (2008)
- [25] G. Isidori and P. Paradisi, *Phys. Lett. B* **639**, 499 (2006).
- [26] Belle Collaboration, K. Ikado, et al. *Phys. Rev. Lett.* **97**, 251802 (2006); BaBar Collaboration, B. Aubert, et al. arXiv:0705.1820, 2007.